

A Higher Order Method for Multiple Zeros of Nonlinear Functions

H. D. VICTORY, Jr. and B. NETA†

Texas Tech University, Department of Mathematics, Lubbock, TX 79409, U.S.A.

(Received April, 1982)

A method of order three for finding multiple zeros of nonlinear functions is developed. The method requires two evaluations of the function and one evaluation of the derivative per step.

KEY WORDS: Non-linear equations, order of convergence, multiple root, iteration

C.R. CATEGORIES: 5.1, 5.15

1. INTRODUCTION

Newton's method for computing a zero ξ of multiplicity m of a nonlinear equation $f(x)=0$ is of order one. There is a well-known modification (see e.g. Rall [4] or Schröder [5, p. 324]) of the method to obtain a second order. This uses a knowledge of the multiplicity.

Here we develop a method of order three. An iteration consists of one Newton substep followed by a substep of "modified" Newton (i.e., using the derivative of f at the first substep instead of the current one). The research for this article was motivated by the research of the second author in [2], where a sixth-order method was derived for simple roots, and required three evaluations of the given function and only one evaluation

†This author was supported by N.S.F. Grant No. CPE8007396 and by the Institute for University Research-Arts and Sciences, Texas Tech University (Professor Arthur Draper, Director).

of the derivative per step. Indeed, the research in this paper represents initial endeavors in extending the analysis for higher order methods in [2-3] to include the case of multiple roots of either nonlinear functions or of Jacobian determinants of nonlinear systems.

Let us recall the definition of order (see e.g. [2]).

DEFINITION Let x_1, x_2, \dots, x_i be a sequence converging to ξ . Let $e_i = x_i - \xi$. If there exist a real number p and a nonzero constant C such that

$$\frac{|e_{i+1}|}{|e_i|^p} \rightarrow C \quad (1)$$

then p is called the order of the sequence.

2. DEVELOPMENT OF THE THIRD-ORDER METHOD

Let $m > 1$ be the multiplicity of the root ξ to be computed, and suppose f has a least $m+2$ derivatives in a neighborhood of ξ . The i th derivative of our function f is denoted by $f^{(i)}$.

Let

$$\left. \begin{aligned} w_n &= x_n - \frac{f(x_n)}{f'(x_n)} \\ x_{n+1} &= w_n - \frac{f(w_n)f(x_n) + Af(w_n)}{f'(x_n)f(x_n) + Bf(w_n)} \end{aligned} \right\} \quad (2)$$

where A, B are arbitrary constants. This is a family of methods which uses Newton's method in the first substep and a Newton-like in the second one. In each step we have to evaluate the function $f(x)$ at the two points x_n, w_n and to evaluate the derivative at one point x_n .

In order to find the order of the method we used MACSYMA (Project MAC's Symbolic MANipulation system, which is a large computer programming system written in LISP and used for performing symbolic as well as numerical mathematical manipulation [1]).

The error expression at w_n is given by

$$\begin{aligned} e(w_n) = w_n - \xi &= \frac{m-1}{m} e_n + \frac{F_2}{m^2(m+1)} e_n^2 \\ &- \left(\frac{F_2^2}{m^3(m+1)} - \frac{2F_3}{m^2(m+1)(m+2)} \right) e_n^3 + \dots, \end{aligned} \quad (3)$$

where

$$F_i = \frac{f^{(i+m-1)}(\xi)}{f^{(m)}(\xi)}, \quad (4)$$

and

$$\varepsilon_n = \vartheta(x_n) = x_n - \xi. \quad (5)$$

For later use we also give the expression for

$$\begin{aligned} f(w_n)[f(x_n) + Af(w_n)] &= \frac{\mu^m(1 + A\mu^m)}{m!^2} \varepsilon_n^{2m} \left\{ 1 + \varepsilon_n \left[\frac{F_2(2m^2 - 2m + 1)}{m(m^2 - 1)} \right. \right. \\ &\quad \left. \left. + \frac{F_2 A \mu^m}{m(m^2 - 1)(1 + A\mu^m)} \right] + \dots \right\}, \end{aligned} \quad (6)$$

and

$$\begin{aligned} f'(x_n)[f(x_n) + Bf(w_n)] &= \frac{B\mu^m + 1}{m!(m-1)!} \varepsilon_n^{2m-1} \left\{ 1 + \varepsilon_n \left[\frac{F_2(2m^2 - m - 1)}{m(m^2 - 1)(1 + B\mu^m)} \right. \right. \\ &\quad \left. \left. + \frac{F_2 B \mu^m(2m^2 - m)}{m(m^2 - 1)(1 + B\mu^m)} \right] + \dots \right\}, \end{aligned} \quad (7)$$

where

$$\mu = \frac{m-1}{m}. \quad (8)$$

The error expression at the point x_{n+1} is given by

$$\begin{aligned} \varepsilon_{n+1} &= \left[\mu - \frac{\mu^m(1 + A\mu^m)}{m(1 + B\mu^m)} \right] \varepsilon_n + \frac{F_2}{m^2(m+1)} \varepsilon_n^2 \\ &+ \frac{\mu^m(1 + A\mu^m)}{m(1 + B\mu^m)} \left[\frac{(m-2)F_2}{m(m^2 - 1)} - \frac{\mu A F_2}{m(m^2 - 1)(1 + A\mu^m)} + \frac{\mu B F_2}{m(m^2 - 1)(1 + B\mu^m)} \right] \varepsilon_n^2 + \dots \end{aligned}$$

The expression for the coefficient for e_n^3 is too long (over 2 pages) to copy from the computer output.

In order to annihilate the coefficient of e_n the parameters A, B should satisfy:

$$\mu = \frac{\mu^m(1 + A\mu^m)}{m(1 + B\mu^m)}. \quad (10)$$

Combining (9) and (10) one has

$$e_{n+1} = \frac{m-1}{m^2(m+1)} F_2 e_n^2 - \frac{F_2}{m^2(m+1)} \frac{m\mu^{1-m} - 1}{m\mu^{1-m}(1 + B\mu^m)} e_n^2 + \dots \quad (11)$$

To annihilate the second order terms one needs

$$B = -\frac{\left(\frac{m}{m-1}\right)^m (m-2)(m-1) + 1}{(m-1)^2}. \quad (12)$$

Substituting this value of B in (10) one obtains

$$A = \left(\frac{m}{m-1}\right)^{2m} - \left(\frac{m}{m-1}\right)^{m+1}. \quad (13)$$

Note that A, B depend on m . The knowledge of m in this case increased the order to three. For a discussion of how to estimate the multiplicity m , see for example Traub [6, pp. 129–130].

3. NUMERICAL EXAMPLES

In our first experiment we obtained the double root $\xi = 1$ of the function

$$f(x) = x^2 - 2x + 1 \quad (14)$$

using our method and the following modified Newton method

$$x_{n+1} = x_n - m \frac{f(x_n)}{f'(x_n)}. \quad (15)$$

Our method obtained the root in 1 iteration using a starting value $x_0 = 0$. The modified Newton method (15) required 2 iterations.

The second experiment was to obtain a triple root at $\xi=1$ of the function

$$f(x)=x^5-8x^4+24x^3-34x^2+23x-6. \quad (16)$$

Again, we used $x_0=0$ as starting value. The convergence of our method was so fast that we were not able to measure the rate of convergence. The results were obtained in quadruple precision on ITEL AS-6 and are summarized in Table I.

TABLE I

Iteration number	x		Absolute Error	
	Newton	This paper	Newton	This paper
0	0.	0.	1.	1.
1	0.7826087	0.9294938	0.2173913	0.0705062
2	0.9816479	0.9999038	0.0183521	0.0000962
3	0.9998356	1.0000000	0.0001644	0.
4	1.0000000	—	0.	—

TABLE II

x_{old}	x_{new}	$ x_{old}-1 ^3$	$ x_{new}-1 $
0.	0.9294938	1.	$0.705062 \cdot 10^{-1}$
0.9294938	0.9999038	$0.350486 \cdot 10^{-3}$	$0.962 \cdot 10^{-4}$
0.9999038	1.0000000	$0.890277 \cdot 10^{-12}$	0.

In Table II we compute $|x_{old}-1|^3$ and $|x_{new}-1|$ for the second problem. Note that our method produced an error which is smaller than the cube of the error at the previous step.

In our next experiment we were able to measure the rate of convergence. The function

$$f(x)=3x^4+8x^3-6x^2-24x+19 \quad (17)$$

has a double root at $\xi=1$. Starting with $x_0=0$ our method used 2/3 the number of iterations required by the modified Newton. The results are summarized in Table III.

TABLE III

Iteration number	x		rate	
	Newton	This paper	Newton	This paper
0	0.	0.	—	—
1	1.583333	0.8904491	—	—
2	1.071987	0.9998828	—	—
3	1.001386	1.000000	1.7500	3.2624
4	1.000001	1.000000	1.9850	—
5	1.000000	—	2.0002	—
6	1.000000	—	—	—

In our last experiment we compare our method to Werner's [7] for approximating a double root at $\xi=0$ of

$$f(x) = x^2 \exp(x). \quad (18)$$

The results are summarized in Table IV.

TABLE IV

Iteration number	x			rate		
	Werner	This paper	Werner	This paper	Werner	This paper
0	0.2	0.2	0.1	—	—	—
1	$0.2160 \cdot 10^{-1}$	$0.1487 \cdot 10^{-2}$	0.1141	2.383	4.045	—
2	$0.2379 \cdot 10^{-3}$	$0.6844 \cdot 10^{-9}$	$0.2904 \cdot 10^{-3}$	2.176	3.241	3.752
3	$0.2830 \cdot 10^{-7}$	—	$0.5103 \cdot 10^{-11}$	2.083	—	3.193

Acknowledgements

The second author wishes to thank Professor George J. Fix for the invitation to Carnegie-Mellon University and for the computer time. Thanks also to Mr. Victor Trutzer for his help with MACSYMA.

References

- [1] R. Bogen, *et al.*, *MACSYMA Reference Manual*, M.I.T., Cambridge, Mass., 1975.
- [2] B. Neta, A sixth-order family of methods for nonlinear equations, *Intern. J. Computer Math.*, Sec. B. 7 (1979), 157-161.
- [3] B. Neta, A new iterative method for the solution of systems of nonlinear equations, *Proc. Conf. Approx. Th. Appl.* (1981), 249-263, (Z. Ziegler, Ed.).

- [4] L. B. Rall, Convergence of the Newton process to multiple solutions, *Numer. Math.*, **9** (1966), 23-37.
- [5] E. Schröder, Über unendlich viele Algorithmen zur Auflösung der Gleichungen, *Math. Ann.* **2** (1970), 317-365.
- [6] J. F. Traub, *Iterative Methods for the Solution of Equations*, Prentice-Hall, Englewood Cliffs, NJ, 1964.
- [7] W. Werner, Some efficient algorithms for the solution of a single nonlinear equation, *Intern. J. Computer Math. Sec. B*, **9** (1981), 141-149.